

ON SOME HADAMARD-TYPE INEQUALITIES FOR CONVEX FUNCTIONS ON A RECTANGULAR BOX

★M. EMIN ÖZDEMİR AND ♦♣AHMET OCAK AKDEMİR

ABSTRACT. In this paper some Hadamard-type inequalities for convex functions of 3-variables on a rectangular box are given. We also define a mapping related to convex functions on a rectangular box.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a < b$. The following inequality;

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}$$

is known in the literature as Hadamard's inequality for convex mappings. The history of this famous integral inequality begins with the papers of Ch. Hermite [18] and J. Hadamard [17]. In recent years there have been many extensions, generalizations, refinements and similar results of Hadamard's inequality, we refer interest of readers to [1]-[16]. On all of these, in [25], Dragomir defined convex functions on the co-ordinates as following;

Let us consider the bidimensional interval $\Delta := [a, b] \times [c, d]$ in $\mathbb{R}^2$ with $a < b$ and $c < d$. A function $f : \Delta \rightarrow \mathbb{R}$ will be called convex on the co-ordinates if the partial mappings $f_y : [a, b] \rightarrow \mathbb{R}$, $f_y(u) = f(u, y)$ and $f_x : [c, d] \rightarrow \mathbb{R}$, $f_x(v) = f(x, v)$, are convex where defined for all $x \in [a, b]$ and $y \in [c, d]$.

Recall that the mapping $f : \Delta \rightarrow \mathbb{R}$ is convex in Δ if the following inequality:

$$f(\lambda x + (1 - \lambda)z, \lambda y + (1 - \lambda)w) \leq \lambda f(x, y) + (1 - \lambda)f(z, w)$$

holds for all $(x, y), (z, w) \in \Delta$ and $\lambda \in [0, 1]$. Therefore, in [25], Dragomir proved the following lemma and established following inequalities for convex functions on the co-ordinates.

Lemma 1. *Every convex mapping $f : \Delta \rightarrow \mathbb{R}$ is convex on the co-ordinates, but the converse is not generally true.*

Date: September 2010.

1991 *Mathematics Subject Classification.* Primary 26D15, Secondary 26A51.

Key words and phrases. Convex, co-ordinated convex, Co-ordinates, Hadamard's inequality

♣Corresponding author.

Theorem 1. Suppose that $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ is convex on the co-ordinates on Δ . Then one has the inequalities:

$$\begin{aligned}
 & f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \\
 & \leq \frac{1}{2} \left[\frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}\right) dx + \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y\right) dy \right] \\
 (1.1) \quad & \leq \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dx dy \\
 & \leq \frac{1}{4} \left[\frac{1}{b-a} \int_a^b f(x, c) dx + \frac{1}{b-a} \int_a^b f(x, d) dx \right. \\
 & \quad \left. + \frac{1}{d-c} \int_c^d f(a, y) dy + \frac{1}{d-c} \int_c^d f(b, y) dy \right] \\
 & \leq \frac{f(a, c) + f(a, d) + f(b, c) + f(b, d)}{4}
 \end{aligned}$$

The above inequalities are sharp.

In [25], Dragomir considered a mapping which closely connected with above inequalities and established main properties of this mapping as following:

Now, for a mapping $f : \Delta := [a, b] \times [c, d] \rightarrow \mathbb{R}$ is convex on the co-ordinates on Δ , we can define the mapping $H : [0, 1]^2 \rightarrow \mathbb{R}$,

$$H(t, s) := \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f\left(tx + (1-t)\frac{a+b}{2}, sy + (1-s)\frac{c+d}{2}\right) dx dy$$

Theorem 2. Suppose that $f : \Delta \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ is convex on the co-ordinates on $\Delta = [a, b] \times [c, d]$. Then:

- (i) The mapping H is convex on the co-ordinates on $[0, 1]^2$.
- (ii) We have the bounds

$$\sup_{(t,s) \in [0,1]^2} H(t, s) = \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y) dx dy = H(1, 1)$$

$$\inf_{(t,s) \in [0,1]^2} H(t, s) = f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) = H(0, 0)$$

- (iii) The mapping H is monotonic nondecreasing on the co-ordinates.

For similar results see the papers [19]-[26].

In this paper, we defined convex functions on a rectangular box and proved a theorem similar to Theorem 1 on a rectangular box, we also established a mapping related to convex functions on a rectangular box.

2. MAIN RESULT

Firstly we can define the convex function on a rectangular box, as following:

Definition 1. Let us consider the rectangular box $G = [a, b] \times [c, d] \times [e, f]$ in $\mathbb{R}^3$. The mapping $f : G \rightarrow \mathbb{R}$ is convex on G if

$$f(tx + (1-t)u, ty + (1-t)v, tz + (1-t)w) \leq tf(x, y, z) + (1-t)f(u, v, w)$$

holds for all $(x, y, z), (u, v, w) \in G$ and $t \in [0, 1]$.

A function $f : G \rightarrow \mathbb{R}$ is convex on G is called co-ordinated convex on G if the partial mappings

$$f_x : [c, d] \times [e, f] \rightarrow \mathbb{R}, \quad f_x(y, z) = f(x, y, z), \quad x \in [a, b]$$

$$f_y : [a, b] \times [e, f] \rightarrow \mathbb{R}, \quad f_y(x, z) = f(x, y, z), \quad y \in [c, d]$$

$$f_z : [a, b] \times [c, d] \rightarrow \mathbb{R}, \quad f_z(x, y) = f(x, y, z), \quad z \in [e, f]$$

are convex for all $(y, z) \in [c, d] \times [e, f]$, $(x, z) \in [a, b] \times [e, f]$, $(x, y) \in [a, b] \times [c, d]$.

In order to prove our main theorem we need the following lemma.

Lemma 2. Every co-ordinated convex mapping $f : G \rightarrow \mathbb{R}$ is convex for triple co-ordinates, where $G = [a, b] \times [c, d] \times [e, f]$.

Proof. Suppose that $f : G = [a, b] \times [c, d] \times [e, f] \rightarrow \mathbb{R}$ is convex on G . Recall the function

$$f_x : [c, d] \times [e, f] \rightarrow \mathbb{R}, \quad f_x(y, z) = f(x, y, z), \quad x \in [a, b]$$

Then for $t \in [0, 1]$ and $(y_1, z_1), (y_2, z_2) \in [c, d] \times [e, f]$, we have

$$\begin{aligned} f_x(ty_1 + (1-t)y_2, tz_1 + (1-t)z_2) &= f(x, ty_1 + (1-t)y_2, tz_1 + (1-t)z_2) \\ &= f(tx + (1-t)x, ty_1 + (1-t)y_2, tz_1 + (1-t)z_2) \\ &\leq tf(x, y_1, z_1) + (1-t)f(x, y_2, z_2) \\ &= tf_x(y_1, z_1) + (1-t)f_x(y_2, z_2) \end{aligned}$$

Therefore $f_x(y, z) = f(x, y, z)$ is convex on $[c, d] \times [e, f]$, for all $x \in [a, b]$. The fact that $f_y : [a, b] \times [e, f] \rightarrow \mathbb{R}$, $f_y(x, z) = f(x, y, z)$ is convex on $[a, b] \times [e, f]$ for all $y \in [c, d]$ and $f_z : [a, b] \times [c, d] \rightarrow \mathbb{R}$, $f_z(x, y) = f(x, y, z)$ is also convex on $[a, b] \times [c, d]$ for all $z \in [e, f]$ goes likewise and we shall omit the details. $\square$

Theorem 3. Suppose that $f : G = [a, b] \times [c, d] \times [e, f] \rightarrow \mathbb{R}$ is convex on G . Then one has the inequalities:

$$\begin{aligned}
(2.1) \quad & \left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2} \right) \\
\leq & \frac{1}{3(b-a)} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx + \frac{1}{3(d-c)} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy \\
& + \frac{1}{3(f-e)} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz \\
\leq & \frac{1}{(b-a)(d-c)(f-e)} \iiint_G f(x, y, z) dy dz dx \\
\leq & \frac{1}{6} \left[\frac{1}{(b-a)(d-c)} \iint_{\Delta_1} f(x, y, e) dy dx + \frac{1}{(b-a)(d-c)} \iint_{\Delta_1} f(x, y, f) dy dx \right. \\
& + \frac{1}{(b-a)(f-e)} \iint_{\Delta_2} f(x, c, z) dz dx + \frac{1}{(b-a)(f-e)} \iint_{\Delta_2} f(x, d, z) dz dx \\
& + \frac{1}{(d-c)(f-e)} \iint_{\Delta_3} f(a, y, z) dz dy + \frac{1}{(d-c)(f-e)} \iint_{\Delta_3} f(b, y, z) dz dy \left. \right] \\
\leq & \frac{f(a, c, e) + f(a, d, e) + f(b, c, e) + f(b, d, e)}{8} \\
& + \frac{f(a, c, f) + f(a, d, f) + f(b, c, f) + f(b, d, f)}{8}
\end{aligned}$$

where $\Delta_1 = [a, b] \times [c, d]$, $\Delta_2 = [a, b] \times [e, f]$ and $\Delta_3 = [c, d] \times [e, f]$.

Proof. Since $f : G = [a, b] \times [c, d] \times [e, f] \rightarrow \mathbb{R}$ is convex on G , we can write the partial mappings

$$g_x : [c, d] \times [e, f] \rightarrow \mathbb{R}, \quad g_x(y, z) = f(x, y, z)$$

$$g_y : [a, b] \times [e, f] \rightarrow \mathbb{R}, \quad g_y(x, z) = f(x, y, z)$$

$$g_z : [a, b] \times [c, d] \rightarrow \mathbb{R}, \quad g_z(x, y) = f(x, y, z)$$

are convex for all $(y, z) \in [c, d] \times [e, f]$, $(x, z) \in [a, b] \times [e, f]$, $(x, y) \in [a, b] \times [c, d]$. Then by the inequality of (1.1), we have

$$\begin{aligned}
& g_x\left(\frac{c+d}{2}, \frac{e+f}{2}\right) \\
\leq & \frac{1}{(d-c)(f-e)} \int_c^d \int_e^f g_x(y, z) dy dz
\end{aligned}$$

That is

$$\begin{aligned}
& f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) \\
& \leq \frac{1}{(d-c)(f-e)} \int_c^d \int_e^f f(x, y, z) dy dz \\
& = \frac{1}{(d-c)(f-e)} \iint_{\Delta_3} f(x, y, z) dy dz
\end{aligned}$$

Integrating this inequality on $[a, b]$ respect to x and dividing both sides of inequalities $(b-a)$, we have

$$\begin{aligned}
(2.2) \quad & \frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx \\
& \leq \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dy dz dx \\
& = \frac{1}{(b-a)(d-c)(f-e)} \iiint_G f(x, y, z) dy dz dx
\end{aligned}$$

By a similar argument, we have

$$\begin{aligned}
(2.3) \quad & \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy \\
& \leq \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dy dz dx \\
& = \frac{1}{(b-a)(d-c)(f-e)} \iiint_G f(x, y, z) dy dz dx
\end{aligned}$$

$$\begin{aligned}
(2.4) \quad & \frac{1}{f-e} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz \\
& \leq \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dy dz dx \\
& = \frac{1}{(b-a)(d-c)(f-e)} \iiint_G f(x, y, z) dy dz dx
\end{aligned}$$

By addition (2.2), (2.3) and (2.4), we get the second inequality of (2.1)

$$\begin{aligned}
& \frac{1}{3(b-a)} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx + \frac{1}{3(d-c)} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy \\
& + \frac{1}{3(f-e)} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz \\
& \leq \frac{1}{(b-a)(d-c)(f-e)} \iiint_G f(x, y, z) dy dz dx
\end{aligned}$$

By the first inequality of (1.1), we have

$$\begin{aligned}
(2.5) \quad & f\left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2}\right) \\
& \leq \frac{1}{2} \left[\frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy + \frac{1}{f-e} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz \right]
\end{aligned}$$

$$\begin{aligned}
(2.6) \quad & f\left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2}\right) \\
& \leq \frac{1}{2} \left[\frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx + \frac{1}{f-e} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz \right]
\end{aligned}$$

$$\begin{aligned}
(2.7) \quad & f\left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2}\right) \\
& \leq \frac{1}{2} \left[\frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx + \frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy \right]
\end{aligned}$$

Summing the inequalities (2.5)-(2.7), we get the first inequality of (2.1)

$$\begin{aligned}
& f\left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2}\right) \\
& \leq \frac{1}{3} \left[\frac{1}{d-c} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy + \frac{1}{b-a} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx \right. \\
& \quad \left. + \frac{1}{f-e} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz \right]
\end{aligned}$$

Finally, by the third and last inequalities of (1.1), we have

$$\begin{aligned}
 (2.8) \quad & \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dx dy dz \\
 & \leq \frac{1}{4} \left[\frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y, e) dy dx + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y, f) dy dx \right. \\
 & \quad \left. + \frac{1}{(b-a)(f-e)} \int_a^b \int_e^f f(x, c, z) dz dx + \frac{1}{(b-a)(f-e)} \int_a^b \int_e^f f(x, d, z) dz dx \right] \\
 & \leq \frac{\int_a^b f(x, c, e) dx + \int_a^b f(x, c, f) dx + \int_a^b f(x, d, e) dx + \int_a^b f(x, d, f) dx}{4(b-a)}
 \end{aligned}$$

$$\begin{aligned}
 (2.9) \quad & \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dx dy dz \\
 & \leq \frac{1}{4} \left[\frac{1}{(b-a)(d-c)} \int_c^d \int_a^b f(x, y, e) dx dy + \frac{1}{(b-a)(d-c)} \int_c^d \int_a^b f(x, y, f) dx dy \right. \\
 & \quad \left. + \frac{1}{(d-c)(f-e)} \int_c^d \int_e^f f(a, y, z) dz dy + \frac{1}{(d-c)(f-e)} \int_c^d \int_e^f f(b, y, z) dz dy \right] \\
 & \leq \frac{\int_c^d f(a, y, e) dy + \int_c^d f(a, y, f) dy + \int_c^d f(b, y, e) dy + \int_c^d f(b, y, f) dy}{4(d-c)}
 \end{aligned}$$

$$\begin{aligned}
 (2.10) \quad & \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dx dy dz \\
 & \leq \frac{1}{4} \left[\frac{1}{(b-a)(f-e)} \int_e^f \int_a^b f(x, c, z) dx dz + \frac{1}{(b-a)(f-e)} \int_e^f \int_a^b f(x, d, z) dx dz \right. \\
 & \quad \left. + \frac{1}{(d-c)(f-e)} \int_e^f \int_c^d f(a, y, z) dy dz + \frac{1}{(d-c)(f-e)} \int_e^f \int_c^d f(b, y, z) dy dz \right] \\
 & \leq \frac{\int_e^f f(a, c, z) dz + \int_e^f f(a, d, z) dz + \int_e^f f(b, c, z) dz + \int_e^f f(b, d, z) dz}{4(f-e)}
 \end{aligned}$$

Summing the inequalities (2.8)-(2.10) and computing the right hand sides of above inequalities, we get the third and last inequalities of (2.1)

$$\begin{aligned}
& \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dx dy dz \\
& \leq \frac{1}{6} \left[\frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y, e) dy dx + \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x, y, f) dy dx \right. \\
& \quad + \frac{1}{(b-a)(f-e)} \int_a^b \int_e^f f(x, c, z) dz dx + \frac{1}{(b-a)(f-e)} \int_a^b \int_e^f f(x, d, z) dz dx \\
& \quad \left. + \frac{1}{(d-c)(f-e)} \int_c^d \int_e^f f(a, y, z) dz dy + \frac{1}{(d-c)(f-e)} \int_c^d \int_e^f f(b, y, z) dz dy \right] \\
& \leq \frac{f(a, c, e) + f(a, d, e) + f(b, c, e) + f(b, d, e)}{4} \\
& \quad + \frac{f(a, c, f) + f(a, d, f) + f(b, c, f) + f(b, d, f)}{4}
\end{aligned}$$

which completes the proof. $\square$

Now, for a mapping $f : G = [a, b] \times [c, d] \times [e, f] \rightarrow \mathbb{R}$ as above, we can define the mapping $H : [0, 1]^3 \rightarrow \mathbb{R}$,

$$\begin{aligned}
H(t, s, r) & : = \frac{1}{(b-a)(d-c)(f-e)} \\
& \quad \times \int_a^b \int_c^d \int_e^f f \left(tx + (1-t)\frac{a+b}{2}, sy + (1-s)\frac{c+d}{2}, rz + (1-r)\frac{e+f}{2} \right) dx dy dz.
\end{aligned}$$

The following theorem contains the properties of this mapping.

Theorem 4. Suppose that $f : G = [a, b] \times [c, d] \times [e, f] \rightarrow \mathbb{R}$ is convex on G , then:

- (i) The mapping H is convex on the co-ordinates on $[0, 1]^3$.
- (ii) We have the bounds

$$\sup_{(t,s,r) \in [0,1]^3} H(t, s, r) = \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dy dz dx = H(1, 1, 1)$$

$$\inf_{(t,s,r) \in [0,1]^3} H(t, s, r) = f \left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2} \right) = H(0, 0, 0)$$

- (iii) The mapping H is monotonic nondecreasing on the co-ordinates.

Proof. (i) Fix $s, r \in [0, 1]$. For all $\alpha, \beta, \gamma \geq 0$ with $\alpha + \beta + \gamma = 1$ and $t_1, t_2, t_3 \in [0, 1]$, we have

$$\begin{aligned}
& H(\alpha t_1 + \beta t_2 + \gamma t_3, s, r) \\
= & \frac{1}{(b-a)(d-c)(f-e)} \\
& \times \int_a^b \int_c^d \int_e^f f \left((\alpha t_1 + \beta t_2 + \gamma t_3)x + (1 - \alpha t_1 + \beta t_2 + \gamma t_3) \frac{a+b}{2} \right. \\
& \left. , sy + (1-s) \frac{c+d}{2}, rz + (1-r) \frac{e+f}{2} \right) dx dy dz \\
= & \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f \left(\alpha \left(t_1 x + (1-t_1) \frac{a+b}{2} \right) \right. \\
& \left. \beta \left(t_2 x + (1-t_2) \frac{a+b}{2} \right) + \gamma \left(t_3 x + (1-t_3) \frac{a+b}{2} \right) , \right. \\
& \left. sy + (1-s) \frac{c+d}{2}, rz + (1-r) \frac{e+f}{2} \right) dx dy dz \\
\leq & \alpha \frac{1}{(b-a)(d-c)(f-e)} \\
& \times \int_a^b \int_c^d \int_e^f f \left(t_1 x + (1-t_1) \frac{a+b}{2}, sy + (1-s) \frac{c+d}{2}, rz + (1-r) \frac{e+f}{2} \right) dx dy dz \\
& + \beta \frac{1}{(b-a)(d-c)(f-e)} \\
& \times \int_a^b \int_c^d \int_e^f f \left(t_2 x + (1-t_2) \frac{a+b}{2}, sy + (1-s) \frac{c+d}{2}, rz + (1-r) \frac{e+f}{2} \right) dx dy dz \\
& + \gamma \frac{1}{(b-a)(d-c)(f-e)} \\
& \times \int_a^b \int_c^d \int_e^f f \left(t_3 x + (1-t_3) \frac{a+b}{2}, sy + (1-s) \frac{c+d}{2}, rz + (1-r) \frac{e+f}{2} \right) dx dy dz \\
= & \alpha H(t_1, s, r) + \beta H(t_2, s, r) + \gamma H(t_3, s, r).
\end{aligned}$$

By a similar way, one can see that

$$H(t, \alpha s_1 + \beta s_2 + \gamma s_3, r) = \alpha H(t, s_1, r) + \beta H(t, s_2, r) + \gamma H(t, s_3, r)$$

and

$$H(t, s, \alpha r_1 + \beta r_2 + \gamma r_3) = \alpha H(t, s, r_1) + \beta H(t, s, r_2) + \gamma H(t, s, r_3)$$

which completes the proof of (i).

(ii) Using convexity of f on the triple co-ordinates on G and Jensen's integral inequality, we get;

$$\begin{aligned}
H(t, s, r) &\geq \frac{1}{(b-a)} \int_a^b f \left(tx + (1-t) \frac{a+b}{2}, \frac{1}{(d-c)} \int_c^d \left[sy + (1-s) \frac{c+d}{2} \right] dy \right. \\
&\quad \left. , \frac{1}{(f-e)} \int_e^f \left[rz + (1-r) \frac{e+f}{2} \right] dz \right) dx \\
&= \frac{1}{(b-a)} \int_a^b f \left(tx + (1-t) \frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2} \right) dx \\
&\geq f \left(\frac{1}{(b-a)} \int_a^b \left[tx + (1-t) \frac{a+b}{2} \right], \frac{c+d}{2}, \frac{e+f}{2} \right) dx \\
&= f \left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2} \right)
\end{aligned}$$

Since H is convex on the co-ordinates, we have

$$\begin{aligned}
&H(t, s, r) \\
&\leq \frac{rst}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dx dy dz + \frac{rs(1-t)}{(d-c)(f-e)} \int_c^d \int_e^f f\left(\frac{a+b}{2}, y, z\right) dy dz \\
&\quad + \frac{rt(1-s)}{(b-a)(f-e)} \int_a^b \int_e^f f\left(x, \frac{c+d}{2}, z\right) dx dz + \frac{st(1-r)}{(b-a)(d-c)} \int_a^b \int_c^d f\left(x, y, \frac{e+f}{2}\right) dx dy \\
&\quad + \frac{t(1-s)(1-r)}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}, \frac{e+f}{2}\right) dx + \frac{s(1-t)(1-r)}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y, \frac{e+f}{2}\right) dy \\
&\quad + \frac{r(1-t)(1-s)}{(f-e)} \int_e^f f\left(\frac{a+b}{2}, \frac{c+d}{2}, z\right) dz + (1-r)(1-t)(1-s) f\left(\frac{a+b}{2}, \frac{c+d}{2}, \frac{e+f}{2}\right)
\end{aligned}$$

By using the inequalities of (2.1), we obtain

$$H(t, s, r) \leq \frac{1}{(b-a)(d-c)(f-e)} \int_a^b \int_c^d \int_e^f f(x, y, z) dx dy dz$$

this completes the proof of (ii).

(iii) By using Hadamard's inequality, one can see that:

$$H(t, s, r) \geq H(0, s, r)$$

for all $(t, s, r) \in [0, 1]^3$. Let $0 \leq t_1 < t_2 \leq 1$. By the convexity of H , for all $(s, r) \in [0, 1]^2$, we have

$$\frac{H(t_2, s, r) - H(t_1, s, r)}{t_2 - t_1} \geq \frac{H(t_1, s, r) - H(0, s, r)}{t_1} \geq 0.$$

Which completes the proof of (iii). $\square$

REFERENCES

- [1] S. S. Dragomir, Two refinements of Hadamard's inequalities, Coll. Pap. of the Fac. of Sci. Kragujevac (Yugoslavia) 11 (1990), 23-26. ZBL No. 729: 26017.
- [2] R. P. Agarwal and S. S. Dragomir, An application of Hayashi's inequality for differentiable functions, Computers Math. Applic., 32 (6) (1996), 95-99.
- [3] S. S. Dragomir, A mapping in connection to Hadamard's inequality, An Ostro. Akad. Wiss. Math. -Natur (Wien) 128 (1991), 17-20. MR 93h: 26032. ZBL No. 747: 26015.
- [4] S. S. Dragomir, On Hadamard's inequality for convex functions, Math. Balkanica 6 (1992), 215-222. MR 934: 26033.
- [5] S. S. Dragomir, A refinement of Hadamard's inequality for isotonic linear functionals, Tamkang J. Math., 24 (1993), 101-106.
- [6] S. S. Dragomir, Two mappings in connection to Hadamard's inequality, J. Math. Anal. Appl. 167 (1992), 49-56. MR 93m: 26038. ZBL No. 758: 26014.
- [7] S. S. Dragomir, D. Barbu and C. Buşe, A probabilistic argument for the convergence of some sequences associated to Hadamard's inequality, Studia Univ. Babeş-Bolyai, Mathematica, 38 (1993), 29-34.
- [8] J. E. Pecaric and S. S. Dragomir, On some integral inequalities for convex functions, Bull. Mat. Inst. Pol. Iasi, 36 (1990), 19-23. MR 93i: 26019. ZBL No. 765: 26008.
- [9] S. S. Dragomir and N. M. Ionescu, Some integral inequalities for differentiable convex functions, Coll. Pap. of the Fac. of Sci. Kragujevac (Yugoslavia) 13 (1992), 49-56. ZBL No. 770: 26009.
- [10] S. S. Dragomir, J. E. Pecaric and J. Sandor, A note on Jensen-Hadamard inequality, L' Anal. Num. Theor. L'Approx. (Romania) 19 (1990), 21-28. MR 93c: 26016. ZBL No. 743: 26016.
- [11] S. S. Dragomir, Some refinements of Hadamard's inequalities, Gaz. Mat. Metod. (Romania) 11 (1990), 189-191.
- [12] S. S. Dragomir, Some integral inequalities for differentiable convex functions, Contributions, Macedonian Acad. of sci. and arts (Scopie) 16 (1992), 77-80.
- [13] S. S. Dragomir and N. M. Ionescu, Some remarks in convex functions, L' Anal. Num. Theor. L'Approx. (Romania) 21 (1992), 31-36. MR 93m: 26023. ZBL No. 770: 26008.
- [14] S. S. Dragomir, D. M. Milosevic and J. Sandor, On some refinements of Hadamard's inequalities and applications, Univ. Beograd, Publ. Elektrotehn. Fak., Ser. Mat., 4 (1993), 3-10.
- [15] J. E. Pecaric and S. S. Dragomir, A generalization of Hadamard's inequality for isotonic linear functionals, Rodovi Math. (Sarajevo) 7 (1991), 103-107. MR 93k: 26026. ZBL No. 738: 26026.
- [16] G. Zabandan, A new refinement of the Hermite-Hadamard inequality for convex functions, Journal of Inequalities in Pure and Applied Mathematics, 10 (2009), 2, Article No: 45.
- [17] J. Hadamard, Etude sur les proprietes des fonctions entieres et en particulier d'une fonction consideree par Riemann, J. Math. Pures et. Appl., 58 (1893), 171-215.
- [18] Ch. Hermite, Sur deux limites d'une integrale definie, Mathesis, 3 (1883), 82.
- [19] M. A. Latif, M. Alomari, On Hadamard-type inequalities for h -convex functions on the co-ordinates, International Journal of Math. Analysis, 3 (2009), no. 33, 1645-1656.
- [20] M. A. Latif, M. Alomari, Hadamard-type inequalities for product two convex functions on the co-ordinates, International Mathematical Forum, 4 (2009), no. 47, 2327-2338.
- [21] M. Alomari, M. Darus, Hadamard-type inequalities for s -convex functions, International Mathematical Forum, 3 (2008), no. 40, 1965-1975.
- [22] M. Alomari, M. Darus, Co-ordinated s -convex function in the first sense with some Hadamard-type inequalities, Int. Journal Contemp. Math. Sciences, 3 (2008), no. 32, 1557-1567.
- [23] M. Alomari, M. Darus, The Hadamard's inequality for s -convex function of 2-variables on the co-ordinates, International Journal of Math. Analysis, 2 (2008), no. 13, 629-638.
- [24] M. E. Özdemir, E. Set, M. Z. Sarıkaya, Some new Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions, Accepted.
- [25] S. S. Dragomir, On the Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, Taiwanese Journal of Mathematics, 5 (2001), no. 4, 775-788.

- [26] D. Y. Hwang, K. L. Tseng and G. S. Yang, Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane, Taiwanese Journal of Mathematics, 11 (2007), 63-73.

★ATATÜRK UNIVERSITY, K. K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, 25240, KAMPUS, ERZURUM, TURKEY

E-mail address: `emos@atauni.edu.tr`

♦AĞRI İBRAHİM ÇEÇEN UNIVERSITY, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS, 04100, AĞRI, TURKEY

E-mail address: `ahmetakdemir@agri.edu.tr`